

---

---

Math 2150 - Homework # 6

Second order linear ODEs - Theory

---

---

1. Use the Wronskian to show that the following functions are linearly independent on  $I$ .

(a)  $f_1(x) = x, f_2(x) = 3x^2, I = (-\infty, \infty)$

(b)  $f_1(x) = \sin(2x), f_2(x) = \sin(x), I = (-\infty, \infty)$

(c)  $f_1(x) = \frac{1}{x}, f_2(x) = x^2, I = (0, \infty)$

---

2. In this problem we will solve

$$x^2y'' - 5xy' + 8y = 24$$

on the interval  $I = (-\infty, \infty)$ .

(a) Show that  $y_h = c_1x^2 + c_2x^4$  is the general solution to the homogeneous equation  $x^2y'' - 5xy' + 8y = 0$ .

(b) Show that  $y_p = 3$  is a particular solution to  $x^2y'' - 5xy' + 8y = 24$ .

(c) Give a formula for the general solution to  $x^2y'' - 5xy' + 8y = 24$ .

(d) Find the solution to the initial-value problem

$$x^2y'' - 5xy' + 8y = 24, \quad y'(1) = 0, \quad y(1) = -1$$

---

3. In this problem we will solve

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 2e^{2x} + 4x - 12$$

on the interval  $I = (-\infty, \infty)$ .

(a) Show that  $y_h = c_1e^{2x} + c_2xe^{2x}$  is the general solution to the homogeneous equation  $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 0$

- (b) Show that  $y_p = x^2e^{2x} + x - 2$  is a particular solution to  $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 2e^{2x} + 4x - 12$ .
- (c) Give a formula for the general solution to  $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 2e^{2x} + 4x - 12$ .
- (d) Find the solution to the initial-value problem

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = 2e^{2x} + 4x - 12, \quad y'(0) = 0, \quad y(0) = 1$$

---

4. In this problem we will solve

$$2x^2y'' + 5xy' + y = x^2 - x$$

on the interval  $I = (0, \infty)$ .

- (a) Show that  $y_h = c_1x^{-1/2} + c_2x^{-1}$  is the general solution to the homogeneous equation  $2x^2y'' + 5xy' + y = 0$ .
- (b) Show that  $y_p = \frac{1}{15}x^2 - \frac{1}{6}x$  is a particular solution to  $2x^2y'' + 5xy' + y = x^2 - x$ .
- (c) Give a formula for the general solution to  $2x^2y'' + 5xy' + y = x^2 - x$ .
- (d) Find the solution to the initial-value problem

$$2x^2y'' + 5xy' + y = x^2 - x, \quad y'(1) = 1, \quad y(1) = 2$$

---