Math 4740 HW 4 Solutions



(ii) We have
$$P(X=5) = 3/6$$

 $P(X=1) = 2/6$
 $P(X=-15) = 1/6$

So we get the following graph:



(iii)

$$E[X] = (\#1)(\frac{2}{6}) + (\#5)(\frac{3}{6})$$

$$Probability \quad \text{Probability you}$$

$$You \quad \text{win } \#1 \quad \text{win } \#5 \quad \text{vin }$$



Let X be the amount won or 1051. $E[X] = (\#1)(\frac{1}{6} + \frac{1}{6}) + (\#5)(\frac{1}{12} + \frac{1}{12} + \frac{1}{12})$ Probability You win #1 $+ (-\#1S)(\frac{5}{12}) = \# \frac{4 + 1S - 75}{12} = -\frac{4}{56} \frac{56}{12}$ Probability you lose #1S $\approx (-\frac{4}{5}4, 67...)$



 $\begin{array}{l} (c) \quad E[\mathbf{X}] = (0)(0.12) + (1)(0.46) \\ + (2)(0.42) \\ \end{array} \\ = 0 + 0.46 + 0.84 = 1.3 \end{array}$

Thus on average 1.3 heads occur on each experiment over the long term.



((a) P(x>0) means the probability that you win Something since I is the amount won or lost. Thus, $P(X > 0) = \frac{18}{38} + \frac{20}{38} \cdot \frac{18}{38} \cdot \frac{18}{38} = \frac{4059}{6859}$ second branch of ≈ 0.5917... top branch uf tree tree leading to Win \$1 leading to win #1 $(b) E[X] = (\#1) \cdot \left[\frac{18}{38} + \frac{20}{38} \cdot \frac{18}{38} \cdot \frac{18}{38} \right]$ Probability you win \$1 + (-\$I). [20.18.20+20.38.20.38.38] Probability you lare \$I + (- \$3) · [20 - 20 - 20] Probability you lose \$3 $= - \$ \frac{39}{361} \approx - \$ 0.11$ So even though the probability that we win is about 59%, We lose on average about \$0.11 per gume.



$$\begin{array}{l} \left(\alpha \right) \\ P(X=10) = \begin{array}{c} 1 \\ 21 \end{array} \end{array} \end{array} \begin{array}{l} \text{this only happens} \\ \text{When You chose} \\ \text{(D) (F) so only} \\ 1 \text{ way to occur} \\ 1 \text{ way to occur} \\ \text{out of } 21. \end{array}$$

(b) $E[x] = (\frac{1}{21})($10) + (\frac{10}{21})($3) + (\frac{10}{21})(-$4]$ = \$0

$$P(X=-4) = \frac{2!}{2!} = \frac{2!}{2!}$$

$$P(X=-4) = \frac{2!}{2!}$$



$$\begin{split} & \underbrace{\mathbb{X}} = a \mod t \mod t \mod t \pmod{t} \\ & = \underbrace{\mathbb{X}} = \left(-\frac{1}{2} 2 \right) \left(\frac{28}{91} \right) + \left(-\frac{1}{91} \right) \left(\frac{16}{91} \right) + \left(\frac{16}{91} \right) + \left(\frac{1}{9} 0 \right) \left(\frac{1}{91} \right) \\ & = \underbrace{\mathbb{X}} = \left(-\frac{1}{2} 2 \right) \left(\frac{28}{91} \right) + \left(-\frac{1}{9} 1 \right) \left(\frac{16}{91} \right) + \left(\frac{1}{9} 0 \right) \left(\frac{1}{91} \right) \\ & + \left(\frac{1}{9} 1 \right) \left(\frac{32}{91} \right) + \left(\frac{1}{9} 2 \right) \left(\frac{8}{91} \right) + \left(\frac{1}{9} 4 \right) \left(\frac{6}{91} \right) \\ & = \underbrace{\mathbb{X}} =$$



$$\begin{aligned} \widehat{(6)} & \text{Let } X = \text{amount won or lost} \\ \overline{(5)} & \overline{(5)} \\$$

game.

(a) Size of sample space
$$|S| = 6^{3}$$

You lose $-\$1$ if none of the dice match
Your number. Thus,
 $p(-1) = P(\$=-1) = \frac{5 \cdot 5 \cdot 5}{6^{3}} = \frac{125}{216}$
You win $\$1$ if exactly one die matches
Your number. Thus,
 $p(1) = P(\And=1) = \frac{1 \cdot 5 \cdot 5 + 5 \cdot 1 \cdot 5 + 5 \cdot 5 \cdot 1}{6^{3}} = \frac{75}{216}$
You win $\$2$ if exactly two dice match
Your number. Thus,
 $p(2) = P(\And=2) = \frac{1 \cdot 1 \cdot 5 + 1 \cdot 5 \cdot 1 + 5 \cdot (-1)}{6^{3}} = \frac{15}{216}$
You win $\$3$ if all the dice match your
number. Thus,
 $p(3) = P(\And=3) = \frac{1 \cdot 1 \cdot 1}{6^{3}} = \frac{1}{6^{3}} = \frac{1}{216}$



(c) $E[X] = (-\#1)(\frac{125}{216}) + (\#1)(\frac{75}{216})$ $+ (\#2)(\frac{15}{216}) + (\#3)(\frac{1}{216})$ $= - \#\frac{17}{216} \approx - \#0.0787$

8 From previous
$$HW_{J}$$
 the probability space is
 $S = \{(H), (T, H), (T, T, H), (T, T, T, H), (T, T, T, T, H), (T, T, T, T, T, H), (T, T, T, T, T, H), ... \}$
The probability tree looks like this:



Let

$$E = \{(T_{1}T_{1}T_{1}H), (T_{1}T_{1}T_{1}T_{1}H), (T_{1}T_{1}T_{1}T_{1}H), ...\}$$
We want P(E).
You could calculate this in two ways.
Method 1
P(E) = P($\{(T_{1}T_{1}T_{1}T_{1}H)\}$ + P($\{(T_{1}T_{1}T_{1}T_{1}H)\}$ + ...
 $= (\frac{1}{2})^{4} + (\frac{1}{2})^{5} + (\frac{1}{2})^{6} + (\frac{1}{2})^{7} + ...$
 $= (\frac{1}{2})^{4} (1 + \frac{1}{2} + (\frac{1}{2})^{2} + (\frac{1}{2})^{3} + ...]$
 $= \frac{1}{16} [\frac{1}{1 - \frac{1}{2}}] = \frac{1}{16} [2] = \frac{1}{8}$
 $1 + x + x^{3} + x^{3} + ... = \frac{1}{1 - x}$
if $-1 < x < 1$
Method 2
 $P(E) = 1 - P(E) = [-P(\{(H), (T, H), (T, T, H)\})]$
 $= 1 - [\frac{1}{2} + (\frac{1}{2})^{2} + (\frac{1}{2})^{3}]$
 $= 1 - [\frac{1}{2} - \frac{1}{4} - \frac{1}{8}] = \frac{8 - 4 - 2 - 1}{8}$

Thus, $P(E) = 1 - P(E) = 1 - \frac{1}{8} = \frac{1}{8}$ Let X be the amount won or lost. Then E[X] = (\$5) P(E) + (-\$1) P(E) $= (\$5) (\frac{1}{8}) + (-\$1) (\frac{2}{8})$ $= -\$\frac{2}{8} = -\0.25

The expected value is negative so in the long run if you did this bet many times you would expect to lose \$0.25 per bet.

(a) $S = \{ (3), (1,3), (2,3), (4,3), (4,3) \}$ (1,1,3), (1,2,3), (1,4,3), (2,1,3),(2,2,31, (2,4,3), (4,1,3), (4,2,3), (4,4,3), ... 7 (1,1,3) (1,3) (3) -19 lhe probability 14 14 (1, 1)tree is (リ2 14 \|4 on the 14 14 14 (1,2) 17 left. 14 (1)The boxed (1,4, 44 1/4 (1, 4)44 elements 44 are S. (2,3) ****/ч Each Vy 4 branch is (2) 14 (2,1) 14 1/4 probability 14 14 (2,2) Yy. You 44 44 (2,2,3) VIY multiply (2,41 44 2,4,3() the probabiliter (4.3) Yu (4,1,3) Yч sce (4,1)44 next VG (4,21 (4) 6 (4,2,3) lu page 44 (4,41 (4,4,3) 119 +

We have $P(\{(3)\}) = \frac{1}{4}$ $P(\{(1,3)\}) = \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^2}$ $P(\{2,3\}) = \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^2}$ $P(\{(4,3)\}) = \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^2}$ $P(\{(1,1,3)\}) = \frac{1}{4} \cdot \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^3}$ $P(\{(1,2,3)\}) = \frac{1}{4} \cdot \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^3}$ $P(\{(1,4,3)\}) = \frac{1}{4} \cdot \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^3}$ $P(\{(z,1,3)\}) = \frac{1}{4} \cdot \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^3}$ $P(\{(2,2,3)\}) = \frac{1}{4} \cdot \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^3}$ $P(\{(2,4,3)\}) = \frac{1}{4} \cdot \frac{1}{4} \cdot \frac{1}{4} = \frac{1}{4^3}$

and so on ...

We check that P is a probability function
by showing that the sum of P over S
is 1. We have
$$\sum_{w \in S} P(\{w\}) = P(\{(1,1,3)\}) + P(\{(2,1,3)\}) + P(\{(2,1,3)\}) + P(\{(2,2,3)\}) + P(\{(2,2,3)\}) + P(\{(2,2,3)\}) + P(\{(2,2,3)\}) + P(\{(2,2,3)\}) + P(\{(2,2,3)\}) + P(\{(2,3,3)\}) + P(\{(3,3,3)\}) + P(\{(3,3,3)\})$$

$$= \frac{1}{4} \left[1 + \frac{3}{4} + \left(\frac{3}{4}\right)^{2} + \left(\frac{3}{4}\right)^{3} + \left(\frac{3}{4}\right)^{4} + \dots \right)$$

$$= \frac{1}{4} \cdot \left[\frac{1}{1 - \frac{3}{4}} \right] = \frac{1}{4} \cdot \left[\frac{1}{\frac{1}{4}} \right] = 1$$

$$\frac{1}{4} \cdot \frac{1}{4} + \frac{$$

(c)
$$B = \{(3), (1,3), (2,3), (4,3), (1,1,3), (1,2,3), (1,4,3), (1,4,3), (2,2,3), (2,4,3), (4,1,3), (4,2,3), (4,2,3), (4,4,3)\}$$

 $P(B) = \frac{1}{4} + 3 \cdot \frac{1}{4^2} + 9 \cdot \frac{1}{4^3} = \frac{16 + 12 + 9}{64}$
 $= \frac{37}{64} \approx 0.578125... \approx 57.8\%$
(d) $X = anount won or lost$
 $E[X] = (\#5)(\frac{37}{64}) + (-\#6)(\frac{27}{64})$
 $\xrightarrow{Probability}_{3 \text{ is rolled}}_{3 \text{ rolls}}$
 $= \frac{\#185 - \#162}{64} = \#\frac{23}{64} \approx \#0.359$
If you can play the game many times then you'd
expect to win on a userage $\#0.36$ per game.
So good to play if you can play many times.