

**California State University – Los Angeles**  
**Department of Mathematics**  
**Master's Degree Comprehensive Examination**  
**Analysis    Fall 2022**  
**Da Silva, Krebs\*, Zhong**

---

Do at least five of the following seven problems. All problems count equally. If you attempt more than five, the best five will be used.

- (1) Write in a fairly soft pencil (number 2) (or in ink if you wish) so that your work will duplicate well. There should be a supply available.
- (2) Write on one side of the paper only.
- (3) Begin each problem on a new page.
- (4) Assemble the problems you hand in in numerical order.

**Exams are graded anonymously, so put your name only where directed and follow any instructions concerning identification code numbers.**

---

**SECTION 1 – Do two (2) problems from this section. If you attempt all three, then the best two will be used for your grade.**

**Fall 2022 #1.** Consider the sequence defined by  $x_0 = 1$  and

$$x_{n+1} = 1 + \frac{1}{x_n}$$

for all integers  $n \geq 0$ .

(a) Show that the sequence satisfies

$$1 \leq x_n \leq 2$$

for all non-negative integers  $n$ .

(b) Prove that  $(x_n)$  has a convergent subsequence  $(x_{n_k})$ . Hint: Use your answer from (a).

**Fall 2022 #2.** Let  $(x_n)_{n=1}^{\infty}$  be a sequence of real numbers.

(a) Define what it means for  $(x_n)_{n=1}^{\infty}$  to be a “Cauchy sequence.”

(b) Use your answer from (a) to prove that if  $(x_n)_{n=1}^{\infty}$  is a Cauchy sequence, then  $\{x_n \mid n \in \mathbb{N}\}$  is bounded. (Here  $\mathbb{N}$  denotes the set of positive integers.)

**Fall 2022 #3.**

Let  $f : D \rightarrow \mathbb{R}$  be a continuous function on an open interval  $D$ . Prove that the function  $f_+ : D \rightarrow \mathbb{R}$  defined by

$$f_+(x) = \max\{f(x), 0\}$$

is continuous.

---

**SECTION 2 – Do three (3) problems from this section. If you attempt more than three, then the best three will be used for your grade.**

**Fall 2022 #4.** Let  $T : C([0, 1]) \rightarrow \mathbb{R}$  be the bounded linear transformation defined by

$$T(f) = \int_0^1 f(x) \, dx.$$

Here  $C([0, 1])$  denotes the space of continuous functions from  $[0, 1]$  to  $\mathbb{R}$ . We endow  $C([0, 1])$  with the  $L^\infty$  norm defined by

$$\|f\|_\infty = \sup\{|f(x)| : 0 \leq x \leq 1\}.$$

- (a) Show that  $\|T\| \leq 1$ .  
 (b) If  $g \in C([0, 1])$  is defined by  $g(x) = 1$ , find  $|T(g)|$ , and use this to compute  $\|T\|$ .

**Fall 2022 #5.** Define

$$\ell^2(\mathbb{N}; \mathbb{R}) := \left\{ (x_n)_{n=1}^\infty \mid x_1, x_2, x_3, \dots \in \mathbb{R} \text{ and } \sum_{n=1}^\infty x_n^2 < \infty \right\}.$$

In other words,  $\ell^2(\mathbb{N}; \mathbb{R})$  is the set of all “square-summable” sequences of real numbers. Recall that  $\ell^2(\mathbb{N}; \mathbb{R})$  is an inner product space with the inner product

$$\langle (x_n)_{n=1}^\infty, (y_n)_{n=1}^\infty \rangle = \sum_{n=1}^\infty x_n y_n.$$

(You may assume without proof that this defines an inner product.)

Define

$$r_n = \begin{cases} 2^{-(n+3)/4} & \text{if } n \text{ is odd} \\ 2^{-(n+2)/4} & \text{if } n \text{ is even} \end{cases}$$

and  $s_n = (-1)^n r_n$ . So

$$(r_n)_{n=1}^{\infty} = (2^{-1}, 2^{-1}, 2^{-3/2}, 2^{-3/2}, 2^{-2}, 2^{-2}, 2^{-5/2}, 2^{-5/2}, \dots), \text{ and}$$

$$(s_n)_{n=1}^{\infty} = (2^{-1}, -2^{-1}, 2^{-3/2}, -2^{-3/2}, 2^{-2}, -2^{-2}, 2^{-5/2}, -2^{-5/2}, \dots).$$

(a) Prove that  $(r_n)_{n=1}^{\infty} \in \ell^2(\mathbb{N}; \mathbb{R})$  and  $(s_n)_{n=1}^{\infty} \in \ell^2(\mathbb{N}; \mathbb{R})$ . (Recall the geometric series formula: If  $x$  is a real number such that  $-1 < x < 1$ , then  $\sum_{n=1}^{\infty} x^n = \frac{x}{1-x}$ .)

(b) Prove that  $\{(r_n)_{n=1}^{\infty}, (s_n)_{n=1}^{\infty}\}$  is an orthonormal family in  $\ell^2(\mathbb{N}; \mathbb{R})$ .

(c) Find real numbers  $a$  and  $b$  so that the quantity  $J(a, b)$  below is as small as possible.

$$J(a, b) = \sum_{n=1}^{\infty} (2^{-n} - ar_n - bs_n)^2.$$

**Fall 2022 #6.** Let  $V$  be a normed vector space over  $\mathbb{F}$ , where  $\mathbb{F}$  may be either the field of real numbers or the field of complex numbers. Let  $A$  be a linear subspace of  $V$ . Let  $C$  be the set of all  $x \in V$  such that there is a sequence  $\{x_n\}_{n=1}^{\infty}$  in  $A$  converging to  $x$ . (In other words,  $C$  is the closure of  $A$ .) Prove that  $C$  is a linear subspace of  $V$ .

**Fall 2022 #7.** Let  $f(x) = x(\pi - x)$  for  $x \in (0, \pi)$ .

(a) Extend the function  $f$  to the interval  $(-\pi, \pi)$  such that it becomes an odd function. Please write down the expression of the extended function  $F(x)$  on  $(-\pi, \pi)$ .

(b) We extend  $F(x)$  from Part (a) to be  $2\pi$ -periodic on  $\mathbb{R}$ . Find the Fourier series for  $F(x)$  in the trigonometric form.

(c) Use the result of Part (b) to find the value of the infinite series

$$1 - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} + \frac{1}{9^3} - \dots$$