

Minimum span of no-hole $(r + 1)$ -distant colorings

Gerard J. Chang* Justie Su-tzu Juan† Daphne Der-Fen Liu‡

May 14, 1998 (Revised April 22, 1999 and September 2000)

Abstract

Given a non-negative integer r , a no-hole $(r + 1)$ -distant coloring, called N_r -coloring, of a graph G is a function that assigns a non-negative integer (color) to each vertex such that the separation of the colors of any pair of adjacent vertices is greater than r , and the set of the colors used must be consecutive. Given r and G , the minimum N_r -span of G , $\text{nsp}_r(G)$, is the minimum difference of the largest and the smallest colors used in an N_r -coloring of G if there exists one; otherwise, define $\text{nsp}_r(G) = \infty$. The values of $\text{nsp}_1(G)$ ($r = 1$) for bipartite graphs are given by Roberts [16]. Given $r \geq 2$, we determine the values of $\text{nsp}_r(G)$ for all bipartite graph with at least $r - 2$ isolated vertices. This leads to complete solutions of $\text{nsp}_2(G)$ for bipartite graphs.

Keywords: Vertex-coloring, no-hole $(r + 1)$ -distant coloring, minimum span, bipartite graphs.

AMS subject classifications: 05C78.

1 Introduction

The T -coloring of graphs models the *channel assignment problem* introduced by Hale [6] in communication networks. In the channel assignment problem, several transmitters and a forbidden set T (called T -set) of non-negative integers with $0 \in T$, are

*Department of Applied Mathematics, National Chiao Tung University, Hsinchu 30050, Taiwan. Email: gjchang@math.nctu.edu.tw. Supported in part by the National Science Council under grant NSC87-2115-M009-007, and the Lee and MTI Center for Networking Research at NCTU.

†Department of Computer Science and Information Engineering, National Chi Nan University, 1, University Road, Puli, Nantou 545, Taiwan. Email: jsjuan@csie.nctu.edu.tw.

‡Corresponding author. Department of Mathematics and Computer Science, California State University, Los Angeles, CA 90032. Email: dliu@calstatela.edu. Supported in part by the National Science Foundation under grant DMS-9805945.

given. We assign a non-negative integral channel to each transmitter under the constraint that if two transmitters interfere, the difference of their channels does not fall within the given T -set. Two transmitters may interfere due to various reasons such as geographic proximity and meteorological factors. To formulate this problem, we construct a graph G such that each vertex represents a transmitter, and two vertices are adjacent if their corresponding transmitters interfere.

Thus, we have the following definition. Given a T -set and a graph G , a T -coloring of G is a function $f : V(G) \rightarrow Z^+ \cup \{0\}$ such that

$$|f(x) - f(y)| \notin T, \text{ if } xy \in E(G).$$

Note that if $T = \{0\}$, then T -coloring is the same as ordinary vertex-coloring.

A *no-hole T -coloring* of a graph G is a T -coloring f of G such that the set $\{f(v) : v \in V(G)\}$ is consecutive (the no-hole assumption). When $T = \{0, 1\}$ and $T = \{0, 1, 2, \dots, r\}$, a no-hole T -coloring is also called an N -coloring [16] and an N_r -coloring (or *no-hole $(r+1)$ -distant coloring*) [17], respectively. That is, an N_r -coloring of a graph G is a vertex coloring $f : V(G) \rightarrow Z^+ \cup \{0\}$ such that the following two conditions are satisfied:

- $|f(x) - f(y)| \geq r + 1$, if $uv \in E(G)$;
- the set $\{f(v) : v \in V(G)\}$ is consecutive.

In terms of efficiency of the usage of the channels (colors), the variable T -span has been considered. The *span* of a T -coloring f is the difference of the largest and the smallest colors used in $f(V)$; the T -span of a graph G , $\text{sp}_T(G)$, is the minimum span among all T -colorings of G .

The T -spans for different families of graphs and for different T -sets have been studied extensively (see [3, 4, 5, 10, 11, 12, 14, 15, 18]). It is known [3, 10] that if T is an *r -initial set*, that is, $T = \{0, 1, 2, \dots, r\} \cup A$ where A is a set of integers without

multiples of $(r + 1)$, then the following holds for all graphs:

$$\text{sp}_T(G) = (\chi(G) - 1)(r + 1), \quad (*)$$

where $\chi(G)$, the *chromatic number* of G , is the minimum number of colors to properly color vertices of G .

It is known [3] and not difficult to learn that for any given T -set and any graph G , a T -coloring always exists. However, a no-hole T -coloring does not always exist. For instance, as $T = \{0, 1\}$, then K_n , complete graph with n vertices, does not have a no-hole T -coloring for any $n \geq 2$.

The minimum span of a no-hole T -coloring for a graph G is denoted by $\text{nsp}_T(G)$. If there does not exist a no-hole T -coloring for G , then $\text{nsp}_T(G) = \infty$. If $T = \{0, 1, 2, \dots, r\}$, denote $\text{nsp}_T(G)$ by $\text{nsp}_r(G)$.

A no-hole T -coloring is also a T -coloring. Hence by (*), a natural lower bound for $\text{nsp}_r(G)$ is $(\chi(G) - 1)(r + 1)$. Roberts [16] and Sakai and Wang [17] studied N -coloring and N_r -coloring, respectively. Among the findings in [16, 17] are the results about the existence of an N -coloring and an N_r -coloring for several families of graphs including paths, cycles, bipartite graphs and 1-unit sphere graphs. The authors also compare the span of such a coloring (if there exists one) with the lower bound $(\chi(G) - 1)(r + 1)$. The N -colorings and N_r -colorings studied in [16, 17] are not necessarily optimal, i.e., the spans are not always the minimum.

This article focuses on the exact values of the minimum N_r -span, $\text{nsp}_r(G)$, especially for bipartite graphs, i.e., graphs with $\chi(G) \leq 2$. In Section 2, we give preliminary results for general graphs. In Section 3, we explore the values of $\text{nsp}_r(G)$ for bipartite graphs. The solutions of $\text{nsp}_1(G)$ for bipartite graphs are given by Roberts [16]. We determine the values of $\text{nsp}_r(G)$ for any bipartite graph G with at least $r - 2$ isolated vertices. This result also leads to a complete description of the values of $\text{nsp}_2(G)$ for all bipartite graphs.

2 Preliminary results

In this section, we present several results about the minimum N_r -span for general graphs. We show a number of upper and lower bounds of $\text{nsp}_r(G)$ for different types of graphs. In order to find the minimum span, without loss of generality, we assume that the color 0 is always used in any N_r -coloring.

If $|V(G)| = n$ and $\text{nsp}_T(G) < \infty$, then by definition, a trivial upper bound for $\text{nsp}_T(G)$ is $n - 1$. On the other hand, any no-hole T -coloring is also a T -coloring, hence we have:

Proposition 1 *For any T -set and any graph G with n vertices, $\text{sp}_T(G) \leq \text{nsp}_T(G)$; and $\text{nsp}_T(G) \leq n - 1$ if $\text{nsp}_T(G) < \infty$.*

Combining Proposition 1 and (*), we have:

Proposition 2 *For any $r \in \mathbb{Z}^+$ and any graph G with chromatic number $\chi(G)$, $(\chi(G) - 1)(r + 1) \leq \text{nsp}_r(G)$.*

With the following result, we show a lower bound of $\text{nsp}_r(G)$ in terms of r and the number of isolated vertices in G .

Theorem 3 *Suppose $r \in \mathbb{Z}^+$ and G is a graph with i isolated vertices, $i \geq 0$, and at least one edge. Then $\text{nsp}_r(G) \geq \max\{2r - i + 1, r + 1\}$.*

Proof. It suffices to show the result when $\text{nsp}_r(G)$ is finite. Because G has at least one edge, $\text{nsp}_r(G) \geq r + 1$. Thus the lemma holds if $i \geq r$.

Suppose $i < r$. Let f be an optimal N_r -coloring of G . By the no-hole assumption of an N_r -coloring, the colors $r, r - 1, \dots, 2, 1, 0$ must be used by some vertices. Since G has only i isolated vertices and $i < r$, there exists a non-isolated vertex u with $r - i \leq f(u) \leq r$. Because u is non-isolated, there exists some vertex v such that

$uv \in E(G)$. Then $f(v) \geq f(u)$, for otherwise $0 \leq f(u) - f(v) \leq r$ contradicting to $uv \in E(G)$. Therefore, we have

$$f(v) \geq f(u) + r + 1 \geq r - i + r + 1 = \max\{2r - i + 1, r + 1\}.$$

This implies $\text{nsp}_r(G) \geq \max\{2r - i + 1, r + 1\}$. □

The *union* of two *vertex-disjoint* graphs G and H , denoted by $G \cup H$, is the graph with vertex set $V(G \cup H) = V(G) \cup V(H)$ and edge set $E(G \cup H) = E(G) \cup E(H)$. For the case in which H has exactly one vertex x , $G \cup H$ is denoted by $G \cup \{x\}$.

The inequality $\text{nsp}_r(G) \leq \text{nsp}_r(G \cup H)$ does not always hold. For instance, if $G = K_2$ then $\text{nsp}_1(G) = \infty$; while $\text{nsp}_1(G \cup \{x\}) = 2$. In the rest of the section, we present several results on unions of graphs.

Theorem 4 *Suppose G is a graph with at least one edge, then $\text{nsp}_{r+1}(G \cup \{x\}) \geq \text{nsp}_r(G) + 1$.*

Proof. It suffices to show the result when $\text{nsp}_{r+1}(G \cup \{x\})$ is finite. Suppose f is an \mathbb{N}_{r+1} -coloring of $G \cup \{x\}$. Define a coloring g on $V(G)$ by:

$$g(v) = \begin{cases} f(v), & \text{if } f(v) < f(x) \text{ or } f(v) = 0; \\ f(v) - 1, & \text{if } f(v) \geq f(x) \text{ and } f(v) > 0. \end{cases}$$

It is straightforward to verify that g is an \mathbb{N}_r -coloring of G and the span of g is one less than the span of f . Therefore, $\text{nsp}_{r+1}(G \cup \{x\}) \geq \text{nsp}_r(G) + 1$. □

Theorem 5 *Suppose G is a graph with $\text{nsp}_r(G) = q(r + 1) + j$, where $q \geq 1$ and $0 \leq j \leq r$, and H is a graph with q vertices. Then $\text{nsp}_{r+1}(G \cup H) \leq \text{nsp}_r(G) + q$.*

Proof. It suffices to show the result when $\text{nsp}_r(G) < \infty$. Let f be an optimal \mathbb{N}_r -coloring of G and $f(V(G)) = \{0, 1, \dots, \text{nsp}_r(G)\}$. Suppose $V(H) = \{x_1, x_2, \dots, x_q\}$. Define a coloring g on $G \cup H$, $g : V(G \cup H) \rightarrow \mathbb{Z}^+ \cup \{0\}$, by

$$g(v) = \begin{cases} \lfloor \frac{(r+2)f(v)}{r+1} \rfloor, & \text{if } v \in V(G); \\ k(r + 2) - 1, & \text{if } v = x_k \in V(H). \end{cases}$$

It is enough to show that g is an N_{r+1} -coloring for $G \cup H$. Because f is onto, so $g(V(G \cup H))$ is a consecutive set, indeed $g(V(G \cup H)) = \{0, 1, 2, \dots, \text{nsp}_r(G) + q\}$. If $uv \in E(G \cup H)$, then either $uv \in E(G)$ or $uv \in E(H)$. If $uv \in E(H)$, then $|g(u) - g(v)| \geq r + 2$. If $uv \in E(G)$, without loss of generality, assume $f(u) > f(v)$. Since $f(u) - f(v) \geq r + 1$, we have $\frac{(r+2)f(u)}{r+1} - \frac{(r+2)f(v)}{r+1} \geq r + 2$, so $g(u) - g(v) \geq r + 2$. Hence g is an N_{r+1} -coloring with span $\text{nsp}_r(G) + q$. This completes the proof. \square

Note that the result in Theorem 5 is not always true if the assumption $\text{nsp}_r(G) = q(r + 1) + j$ does not hold. For instance, let $G = K_2 \cup rK_1$ and $H = K_3$, then $\text{nsp}_r(G) = r + 1$ for any r . However, $\text{nsp}_{r+1}(G \cup H) = \infty$ for any $r \geq 4$.

Corollary 6 *If G is a graph with $r + 1 \leq \text{nsp}_r(G) \leq 2r + 1$, then $\text{nsp}_{r+1}(G \cup \{x\}) = \text{nsp}_r(G) + 1$.*

Proof. The corollary follows from Theorems 4 and 5. \square

Consider the graph G in Figure 1. According to Theorem 3, $\text{nsp}_1(G) \geq 3$ and so the labeling in the figure gives that $\text{nsp}_1(G) = 3$. According to Corollary 6, we have $\text{nsp}_2(G \cup \{x\}) = \text{nsp}_1(G) + 1 = 4$.




Figure 1: Optimal N -coloring for G and optimal N_2 -coloring for $G \cup \{x\}$.

3 Main results

In this section, we explore the minimum N_r -span for bipartite graphs. It turns out that the number of isolated vertices in a bipartite graph plays a key role for this problem. We give the values of $\text{nsp}_r(G)$ for all bipartite graphs G with at least $r - 2$ isolated vertices. This result leads to complete solutions of $\text{nsp}_2(G)$ for all bipartite graphs G .

In this section, a bipartite graph is conventionally denoted by $G = (A, B, I, E)$, where I is the set of all isolated vertices and (A, B) is a *bipartition* of all non-isolated vertices such that each edge in G has one end in A and the other in B . A vertex v is called an A -, B - or I -vertex if $x \in A, B$ or I , respectively.

The *bipartite-complement* \widehat{G} of a bipartite graph $G = (A, B, I, E)$ with $E \neq \emptyset$ is the bipartite graph \widehat{G} with vertex set $V(\widehat{G}) = A \cup B$ and edge set

$$E(\widehat{G}) = \{ab : a \in A, b \in B, ab \notin E\}.$$

Note that the set of isolated vertices in \widehat{G} is not specified in the notation. Moreover, we shall denote B' the set of all B -vertices not adjacent to any A -vertex in \widehat{G} . If G is a bipartite graph, then \widehat{G} is a subgraph of G^c , the *complement* of G (i.e., $V(G^c) = V(G)$ and $E(G^c) = \{uv : u \neq v \text{ and } uv \notin E(G)\}$).

The N_1 -coloring for bipartite graphs has been studied by Roberts [16]. Although the concept of the *minimum* N_1 -span was not introduced explicitly in [16], the following theorem, which completely determines the values of $\text{nsp}_1(G)$ for bipartite graphs, can be generated from [16].

Theorem 7 (Roberts [16]) *If $G = (A, B, I, E)$ is a bipartite graph with $E(G) \neq \emptyset$, then*

$$\text{nsp}_1(G) = \begin{cases} 2, & \text{if } |I| \geq 1; \\ 3, & \text{if } |I| = 0 \text{ and } E(\widehat{G}) \neq \emptyset; \\ \infty, & \text{if } |I| = 0 \text{ and } E(\widehat{G}) = \emptyset. \end{cases}$$

As examples to Theorem 7, consider the graphs G_1 and G_2 in Figure 2. As $|I| \geq 1$ for G_1 , we have $\text{nsp}_1(G_1) = 2$. For G_2 , the facts $|I| = 0$ and $E(\hat{G}) \neq \emptyset$ imply $\text{nsp}_2(G_2) = 3$.




Figure 2: Two examples of optimal N-colorings for bipartite graphs.

Sakai and Wang [17] characterize the existence of an N_r -coloring by using the Hamiltonian r -path. The d -path on n vertices, v_1, v_2, \dots, v_n , has the edge set $\{v_i v_j : 1 \leq |i - j| \leq d\}$. Figure 3 shows a 2-path with 7 vertices. A 1-path on n vertices is an ordinary path denoted as P_n . A *Hamiltonian d -path* of a graph G is a d -path covering each vertex of G exactly once.




Figure 3: A 2-path with 7 vertices.

Theorem 8 (Sakai and Wang [17]) G has an N_r -coloring if and only if G^c has a Hamiltonian r -path. Indeed, if f is an N_r -coloring such that $f(v_1) \leq f(v_2) \leq \dots \leq f(v_n)$, then v_1, v_2, \dots, v_n is a Hamiltonian r -path in G^c .

If the lower bound of $\text{nsp}_r(G)$ in Theorem 3 is attained by some graph G , according to Proposition 2, G must be bipartite. Such graphs do exist. In the next theorem, we show a sufficient condition for such graphs.

Theorem 9 *Suppose $G = (A, B, I, E)$ is a bipartite graph with at least one edge. If $|I| \geq r$, then $\text{nsp}_r(G) = r + 1$; if $|I| \leq r - 1$ and there exist $\{a_1, a_2, \dots, a_{r-|I|}\} \subseteq A$ and $\{b_1, b_2, \dots, b_{r-|I|}\} \subseteq B$ such that $a_j b_k \notin E(G)$ for $j + k \geq r - |I| + 1$, then $\text{nsp}_r(G) = 2r - |I| + 1$.*

Proof. It is obvious that $\text{nsp}_r(G) \geq r + 1$, since $E(G) \neq \emptyset$.

If $|I| \geq r$, coloring A -vertices with 0, B -vertices with $r + 1$, and I -vertices with $1, 2, \dots, r$ gives an N_r -coloring. So, $\text{nsp}_r(G) = r + 1$.

If $|I| \leq r - 1$, by Theorem 3, $\text{nsp}_r(G) \geq 2r - |I| + 1$. Hence, it suffices to find an N_r -coloring with span at most $2r - |I| + 1$. Define a coloring by the following four steps:

- (1) color $a_1, a_2, \dots, a_{r-|I|}$ with $1, 2, \dots, r - |I|$, respectively;
- (2) color I -vertices with $r - |I| + 1, r - |I| + 2, \dots, r$;
- (3) color $b_{r-|I|}, b_{r-|I|-1}, \dots, b_1$ with $r + 1, r + 2, \dots, 2r - |I|$, respectively; and
- (4) color all the remaining A -vertices with 0 and B -vertices with $2r - |I| + 1$.

By the assumption that $a_j b_k \notin E(G)$ for $j + k \geq r - |I| + 1$, it is easy to verify that the coloring defined above is an N_r -coloring with span at most $2r - |I| + 1$. \square

Corollary 10 *Let $G = (A, B, I, E)$ be a bipartite graph with at least one edge.*

- (a) *If $|I| \leq r - 1$ and $E(\widehat{G}) = \emptyset$, then $\text{nsp}_r(G) = \infty$.*
- (b) *If $|I| = r - 1$, then $\text{nsp}_r(G) = r + 2$ iff $E(\widehat{G}) \neq \emptyset$.*
- (c) *If $|I| = r - 2$ and there exists a P_4 in \widehat{G} , then $\text{nsp}_r(G) = r + 3$.*

Proof. We only need to show (a), since (b) and (c) follow from Theorem 9.

Suppose $|I| \leq r - 1$ and $E(\widehat{G}) = \emptyset$. Then, $G - I$ is a complete bipartite graph $K_{|A|,|B|}$. Combining this with the assumption that $|I| \leq r - 1$, G does not admit any N_r -coloring, so $\text{nsp}_r(G) = \infty$. \square

Combining Theorem 9 and Corollary 10 (b), the values of $\text{nsp}_r(G)$ for bipartite graphs with at least $r - 1$ isolated vertices are settled. In the rest of the article, we shall focus on the N_r -coloring for bipartite graphs $G = (A, B, I, E)$ with at most $r - 2$ isolated vertices. By Corollary 10 (a), we may assume $2 \leq |A| \leq |B|$. In the rest of the section, we search for the exact value of $\text{nsp}_r(G)$ to complete the case as $|I| = r - 2$. By Corollary 10 (c), it suffices to consider the case that \widehat{G} contains no P_4 . We first show a lemma which is a key to settle this problem.

For any real number x , denote $\max\{x, 0\}$ by x^+ . For any two integers a and b , $a \leq b$, let $[a, b]$ denote the set $\{a, a + 1, a + 2, \dots, b\}$.

Lemma 11 *Let $G = (A, B, I, E)$ be a bipartite graph with $2 \leq m = |A| \leq |B|$, $|I| \leq r - 2$, and \widehat{G} contains no P_4 . If $\text{nsp}_r(G) < \infty$, then the following are all true:*

- (a) *In the graph \widehat{G} , every B -vertex is adjacent to at most one A -vertex.*
- (b) *There exist an arrangement $\Pi = (A_1, A_2, \dots, A_m)$ of A and non-negative integers $d_1 = 0, c_1, d_2, c_2, d_3, \dots, d_m, c_m = 0$ such that $\deg_{\widehat{G}}(A_k) = d_k + c_k$ for $1 \leq k \leq m$ and $|I| \geq q(\Pi) := \sum_{k=1}^{m-1} q_k$, where $q_k = \max\{(r - c_k)^+, (r - d_{k+1})^+\}$.*
- (c) $\text{nsp}_r(G) \geq (m - 1)(2r + 1) - |I|$.
- (d) *If $B' \neq \emptyset$, then $|I| - q(\Pi) \geq q'(\Pi) := \min_{1 \leq k \leq m-1} q'_k$, where $q'_k = \min\{(r - c_k)^+, (r - d_{k+1})^+\}$.*
- (e) *If $B' \neq \emptyset$, then $\text{nsp}_r(G) \geq \max\{2r + 2, (m - 1)(2r + 1) - |I| + s(\Pi) + 1\}$, where $s(\Pi) = \min_{1 \leq k \leq m-1} \{q_k : q'_k \leq |I| - q(\Pi)\}$.*

Proof. Suppose f is an optimal N_r -coloring for G . According to Theorem 8, G^c has a Hamiltonian r -path $P = v_1, v_2, \dots, v_{|V(G)|}$ with $0 = f(v_1) \leq f(v_2) \leq \dots \leq f(v_{|V(G)|})$. Without loss of generality, we assume the order of A -vertices on the r -path P is $\Pi = (A_1, A_2, \dots, A_m)$. We call this an *arrangement* of A . Hence $f(A_1) \leq f(A_2) \leq \dots \leq f(A_m)$.

On P , let an A - (or B -) *run* be a maximal interval of consecutive $A \cup I$ - (or $B \cup I$ -) vertices, starting and ending with A - (or B -) vertices. Note that there may exist some I -vertices within one run or between two consecutive runs; and the runs are alternating between A and B .

It is impossible to have two consecutive runs with at least two vertices in each. For if it is possible, then there exist $x, y \in A$ and $z, w \in B$ whose order in P is (x, y, z, w) , and the vertices between x and w , other than y and z , are I -vertices. Because $|I| \leq r - 2$, $(x - z - y - w)$ forms a P_4 in \widehat{G} , a contradiction.

Analogously it is impossible to have two consecutive singleton runs (except possibly the first run and the last run). For if it is possible, then we get a P_4 in \widehat{G} by connecting the two consecutive singleton A -run and B -run with the B -vertex and A -vertex preceding and after them.

We conclude that either all A -runs or all B -runs are singletons. As $|A| \leq |B|$, all A -runs are singletons and each B -run (except possibly the first run and/or the last run) contains at least two vertices. So between any A_k and A_{k+1} on P , there are only B - or I -vertices. Since $|I| \leq r - 2$ and P is an Hamiltonian r -path in G^c , there exist at least two B -vertices between A_k and A_{k+1} that are adjacent to A_k .

To prove (a), suppose to the contrary that there exists $v \in B$ such that $vA_k, vA_\ell \in E(\widehat{G})$ for some $k < \ell$. Then between A_k and A_ℓ on P there exists $u \in B - \{v\}$ adjacent to A_k in \widehat{G} . Thus $(u - A_k - v - A_\ell)$ forms a P_4 in \widehat{G} , a contradiction. This proves (a).

Claim: For all $1 \leq k \leq m - 1$, we have $f(A_{k+1}) - f(A_k) \geq r + 2$.

Proof) Suppose $f(A_{k+1}) - f(A_k) \leq r + 1$ for some k . Then the B -vertices between A_k and A_{k+1} on P are adjacent to both A_k and A_{k+1} in \widehat{G} , contradicting (a). \square

Note that if $A_1 = v_i$, then $P' = v_i, v_{i-1}, \dots, v_2, v_1, v_{i+1}, v_{i+2}, \dots, v_{|V(G)|}$ is also a Hamiltonian r -path in G^c , or equivalently, f' defined by $f'(v_j) = f(v_1) + f(v_i) - f(v_j)$ for $1 \leq j \leq i$ and $f'(v_j) = f(v_j)$ for $i < j \leq |V(G)|$ is also an optimal N_r -coloring of G . So, without loss of generality, we may assume $A_1 = v_1$. Similarly, we may assume that $A_m = v_{|V(G)|}$. Put

$$\begin{aligned} D_1 &:= \{y \in B : yA_1 \in E(\widehat{G}) \text{ and } f(y) < f(A_1)\} \text{ and } d_1 := |D_1|; \\ C_1 &:= \{x \in B : xA_1 \in E(\widehat{G}) \text{ and } f(A_1) \leq f(x)\} \text{ and } c_1 := |C_1|; \\ D_k &:= \{y \in B : yA_k \in E(\widehat{G}) \text{ and } f(y) \leq f(A_k)\} \text{ and } d_k := |D_k| \text{ for } 2 \leq k \leq m; \\ C_k &:= \{x \in B : xA_k \in E(\widehat{G}) \text{ and } f(A_k) < f(x)\} \text{ and } c_k := |C_k| \text{ for } 2 \leq k \leq m; \\ I_k &:= \{z \in I : f(A_k) < f(z) < f(A_{k+1})\} \text{ and } i_k := |I_k| \text{ for } 1 \leq k \leq m-1; \\ I'_k &:= \{z \in I : f(A_k) < f(z) \leq f(A_k) + r\} \text{ and } i'_k := |I'_k| \text{ for } 1 \leq k \leq m-1; \\ I''_k &:= \{z \in I : f(A_{k+1}) - r \leq f(z) < f(A_{k+1})\} \text{ and } i''_k := |I''_k| \text{ for } 1 \leq k \leq m-1. \end{aligned}$$

Then $d_1 = c_m = 0$ and $\deg_{\widehat{G}}(A_k) = d_k + c_k$ for $1 \leq k \leq m$. By (a), the C_i 's and D_j 's are all disjoint. By the Claim, for any $1 \leq k \leq m$, $I'_k \cup I''_k \subseteq I_k$ (while I'_k and I''_k are not necessarily disjoint). Furthermore, it is clear that for any $1 \leq k \leq m-1$, $f^{-1}[f(A_k)+1, f(A_k)+r] \subseteq C_k \cup I'_k$, since if $f(A_k) < f(x) \leq f(A_k)+r$, then $x \in C_k \cup I'_k$. Similarly, $f^{-1}[f(A_{k+1})-r, f(A_{k+1})-1] \subseteq D_{k+1} \cup I''_k$. Hence we have $c_k + i'_k \geq r$ and $d_{k+1} + i''_k \geq r$, implying that $i_k \geq \max\{i'_k, i''_k\} \geq \max\{(r - c_k)^+, (r - d_{k+1})^+\} = q_k$ for $1 \leq k \leq m-1$. Therefore,

$$|I| \geq \sum_{k=1}^{m-1} i_k \geq \sum_{k=1}^{m-1} q_k = q(\Pi). \quad (**)$$

This completes the proof of (b).

Now we have $f^{-1}[f(A_k) + 1, f(A_k) + r] \subseteq C_k \cup I'_k \subseteq C_k \cup I_k$ and $f^{-1}[f(A_{k+1}) - r, f(A_{k+1}) - 1] \subseteq D_{k+1} \cup I''_k \subseteq D_{k+1} \cup I_k$. Because $C_k \cap D_{k+1} = \emptyset$, at least $r - i_k$ colors of $[f(A_{k+1}) - r, f(A_{k+1}) - 1]$ are not in $[f(A_k) + 1, f(A_k) + r]$. Thus $f(A_{k+1}) - f(A_k) \geq$

$r + (r - i_k) + 1 = 2r + 1 - i_k$ for $1 \leq k \leq m - 1$. Summing up, we get (c): $\text{nsp}_r(G) \geq f(A_m) - f(A_1) \geq (m - 1)(2r + 1) - |I|$.

Now consider the case that $B' \neq \emptyset$, i.e., there exists some $w \in B$ such that $wA_k \notin E(\widehat{G})$ for all $1 \leq k \leq m$. Hence $|f(w) - f(A_k)| \geq r + 1$, for all $1 \leq k \leq m$. Assume $f(A_p) < f(w) < f(A_{p+1})$ for some $1 \leq p \leq m - 1$. Then $f(A_{p+1}) - f(A_p) \geq 2r + 2$, so $I'_p \cap I''_p = \emptyset$, implying that $i_p \geq i'_p + i''_p \geq (r - c_p)^+ + (r - d_{p+1})^+ = q_p + q'_p$. Replacing $i_p \geq q_p + q'_p$ to the last summation in (**), we get $|I| \geq q(\Pi) + q'_p \geq q(\Pi) + q'(\Pi)$. This proves (d).

Because $f(A_{p+1}) - f(A_p) \geq 2r + 2 \geq 2r + 1 - i_p + q_p + 1$, we have, from the first inequality, $\text{nsp}_r(G) \geq f(A_{p+1}) - f(A_p) \geq 2r + 2$. Using the second inequality, similar to the proof of (c), one can get $\text{nsp}_r(G) \geq (m - 1)(2r + 1) - |I| + q_p + 1 \geq (m - 1)(2r + 1) - |I| + s(\Pi) + 1$. This proves (e). \square

In the next result, we complete the solution of $\text{nsp}_r(G)$ for bipartite graphs $G = (A, B, I, E)$ with $|I| = r - 2$. Let $s(G) = \min s(\Pi)$, where Π runs over all arrangements of A satisfying Lemma 11 (b) and (d).

Theorem 12 *Suppose $G = (A, B, I, E)$ is a bipartite graph with $2 \leq m = |A| \leq |B|$, $0 \leq |I| = r - 2$, and \widehat{G} has no P_4 . Then, $\text{nsp}_r(G) < \infty$ if and only if \widehat{G} satisfies Lemma 11 (a), (b) and (d). In this case,*

$$\text{nsp}_r(G) = \begin{cases} (2r + 1)(m - 1) - r + 2, & \text{if } B' = \emptyset; \\ 2r + 2, & \text{if } B' \neq \emptyset \text{ and } m = 2; \\ (2r + 1)(m - 1) - r + s(G) + 3, & \text{if } B' \neq \emptyset \text{ and } m \geq 3. \end{cases}$$

Proof. The necessity follows from Lemma 11. For the sufficiency, suppose $\Pi = (A_1, A_2, \dots, A_m)$ is an arrangement of A satisfying Lemma 11 (a), (b) and (d). Moreover, assume $s(\Pi) = s(G)$ when $B' \neq \emptyset$.

By Lemma 11 (a), any two A -vertices have disjoint sets of neighbors in \widehat{G} . Then by Lemma 11 (b), we can label the neighbors of A_k in \widehat{G} by $C_{k,1}, C_{k,2}, \dots, C_{k,c_k}$ and $D_{k,1}, D_{k,2}, \dots, D_{k,d_{k+1}}$, respectively, for $1 \leq k \leq m$. In addition, since $|I| \geq \sum_{k=1}^{m-1} q_k$, there exist distinct I -vertices $I_{k,1}, I_{k,2}, \dots, I_{k,q_k}$ for all k .

We shall complete the proof by considering the three cases.

Case 1. $B' = \emptyset$. That is, B is the union of all the C - and D -vertices. It suffices to find an N_r -coloring of G with span $(2r+1)(m-1) - r + 2$. (Then we not only prove that $N_r(G) < \infty$, and also confirm that the span is optimal by Lemma 11(c).) We first replace q_{m-1} by $|I| - \sum_{j=1}^{m-2} q_j$. Then $q_{m-1} \geq \max\{(r - c_{m-1})^+, (r - d_m)^+\}$ and $|I| = \sum_{j=1}^{m-1} q_j$. Indeed, letting B represent the C - and D -vertices and I for I -vertices (without indicating the indices), we can line up all vertices of G as an Hamiltonian r -path in G^c as:

$$P = A_1 \underbrace{BB \cdots B}_{c_1} \underbrace{II \cdots I}_{q_1} \underbrace{BB \cdots B}_{d_2} A_2 \cdots A_{m-1} \underbrace{BB \cdots B}_{c_{m-1}} \underbrace{II \cdots I}_{q_{m-1}} \underbrace{BB \cdots B}_{d_m} A_m.$$

Note that $d_1 = c_m = 0$. Define a coloring on G by the following three steps (the idea is to use each I -vertex to reduce the span by 1).

- (1) A -vertices: $f(A_1) = 0$ and $f(A_{k+1}) = f(A_k) + 2r + 1 - q_k$ for $1 \leq k \leq m - 1$.
- (2) B -vertices: for all $1 \leq k \leq m - 1$,

$$f(C_{k,j}) = \begin{cases} f(A_k) + j, & \text{for } 1 \leq j \leq r - q_k - 1; \\ f(A_k) + r - q_k, & \text{for } r - q_k \leq j \leq c_k, \end{cases}$$

$$f(D_{k+1,j}) = \begin{cases} f(A_k) + r + j, & \text{for } 1 \leq j \leq r - q_k - 1; \\ f(A_k) + 2r - q_k, & \text{for } r - q_k \leq j \leq d_{k+1}. \end{cases}$$

- (3) I -vertices: $f(I_{k,j}) = f(A_k) + r - q_k + j$ for all $q_k > 0$ and $1 \leq j \leq q_k$.

One can easily verify that f is an N_r -coloring for G with span $(2r+1)(m-1) - |I| = (2r+1)(m-1) - r + 2$.

Case 2. $B' \neq \emptyset$ and $m = 2$. Similar to Case 1, by Lemma 11 (e), it suffies to find an N_r -coloring of G with span $\text{nsp}_r(G) = 2r + 2$. Define a coloring by $f(A_1) = 0$, $f(A_2) = 2r + 2$ and $f(z) = r + 1$ for all vertices z in B' . Since $|I| \geq q(\Pi) + q'(\Pi) = q_1 + q'_1 = (r - c_1)^+ + (r - d_2)^+$, there are enough I -vertices to use the colors between 0 and $2r + 2$. Thus one can verify that this is an N_r -coloring of G with span $2r + 2$.

Case 3. $B' \neq \emptyset$ and $m \geq 3$. Again, by Lemma 11 (e), it suffices to find an N_r -coloring with span $(2r + 1)(m - 1) - |I| + s(G) + 1$. Suppose $s(\Pi) = q_p$ for some $1 \leq p \leq m - 1$ with $q'_p \leq |I| - q(\Pi)$. As before, we replace q_i by $q_i + |I| - q(\Pi) - q'_p$ for some $i \neq p$. Then $|I| = q_1 + q_2 + \dots + q_{p-1} + (r - c_p)^+ + (r - d_{p+1})^+ + q_{p+1} + \dots + q_{m-1}$. All the C -, D - and I -vertices are labeled the same as before, except the I -vertices between A_p and A_{p+1} are labeled as $I'_{p,1}, I'_{p,2}, \dots, I'_{p,(r-c_p)^+}, I''_{p,1}, I''_{p,2}, \dots, I''_{p,(r-d_{p+1})^+}$. Apply the same three-step coloring method used for the Case 1, except the colors for the vertices between A_p and A_{p+1} are defined by: $f(I'_{p,j}) = f(A_p) + r - (r - c_p)^+ + j$ for $1 \leq j \leq (r - c_p)^+$; $f(w) = f(A_p) + r + 1$ for all $w \in B'$; $f(I''_{p,j}) = f(A_p) + r + 1 + j$ for $1 \leq j \leq (r - d_{p+1})^+$; $f(A_{p+1}) = f(A_p) + 2r + 2$; and

$$f(C_{p,j}) = \begin{cases} f(A_p) + j, & \text{for } 1 \leq j \leq r - (r - c_p)^+ - 1, \\ f(A_p) + r - (r - c_p)^+, & \text{for } r - (r - c_p)^+ \leq j \leq c_p; \end{cases}$$

$$f(D_{k,j}) = \begin{cases} f(A_p) + r + 1 + (r - d_{p+1})^+ + j, & \text{for } 1 \leq j \leq r - (r - d_{p+1})^+ - 1, \\ f(A_p) + 2r + 1, & \text{for } r - (r - d_{p+1})^+ \leq j \leq d_{p+1}. \end{cases}$$

This gives an N_r -coloring for G with span $(2r + 1)(m - 1) - |I| + s(G) + 1 = (2r + 1)(m - 1) - r + s(G) + 3$. \square

Based on Lemma 11, using a similar process in the proof of Theorem 12, we can also completely settle the case that $I = \emptyset$ and $r \geq 2$. In this case, Lemma 11 (b) means that $q_k = 0$ for all k , or equivalently, that \widehat{G} has two A -vertices of degree at least r and the rest $(m - 2)$ A -vertices of degree at least $2r$. Furthermore, Lemma 11 (d) holds automatically, and $s(\Pi) = 0$. This implies that the lower bound in Lemma 11 (e) is simply $(m - 1)(2r + 1) + 1$. Hence, the same labeling procedure used in Theorem 12 gives the following result.

Theorem 13 *Let $G = (A, B, I, E)$ be a bipartite graph with $2 \leq m = |A| \leq |B|$, $I = \emptyset$, and \widehat{G} contains no P_4 . If $r \geq 2$, then $\text{nsp}_r(G) < \infty$ if and only if Lemma 11 (a) holds and \widehat{G} has two A -vertices of degree at least r and the other $(m - 2)$*

A -vertices of degree at least $2r$. In this case,

$$\text{nsp}_r(G) = \begin{cases} (2r+1)(m-1), & \text{if } B' = \emptyset; \\ (2r+1)(m-1) + 1, & \text{if } B' \neq \emptyset. \end{cases}$$

By Corollary 10 and Theorems 9 and 13, we obtain the complete solutions of $\text{nsp}_2(G)$ for bipartite graphs.


Theorem 14 *If $G = (A, B, I, E)$ is a bipartite graph with at least one edge and $1 \leq m = |A| \leq |B|$, then*

$$\text{nsp}_2(G) = \begin{cases} 3, & \text{if } |I| \geq 2; \\ 4, & \text{if } |I| = 1 \text{ and } E(\widehat{G}) \neq \emptyset; \\ 5, & \text{if } |I| = 0 \text{ and } \widehat{G} \text{ has a } P_4; \\ 5m - 5, & \text{if } |I| = 0, B' = \emptyset \text{ and } \widehat{G} \text{ is a disjoint union of } m \\ & \text{stars, centered at } A \text{ except two of the stars have} \\ & \text{at least 2 edges, each star has at least 4 edges;} \\ 5m - 4, & \text{same as the above, except } B' \neq \emptyset; \\ \infty, & \text{other than any of the above.} \end{cases}$$


Figure 4 shows examples of Theorem 14.

Remark. This article is aimed at computing the values of $\text{nsp}_T(G)$ for bipartite graphs when $T = \{0, 1, \dots, r\}$. Another article by the same authors [1] deals with the values of $\text{nsp}_T(G)$ for unit-interval graphs when $T = \{0, 1\}$. The no-hole T -colorings for some other T -sets and different families of graphs were studied by Liu and Yeh [13]. It was proved [13] that if T is r -initial or $T = [a, b]$, $1 \leq a \leq b$, then for any large n , there exists some graph on n vertices such that $\text{nsp}_T(G)$ equals the upper bound $n - 1$.

Acknowledgment. The authors are grateful to the two anonymous referees for valuable comments.



G_1 : example for case 1 G_2 : example for case 2 G_3 : example for case 3
 $\text{nsp}_2(G_1) = 3$ $\text{nsp}_2(G_2) = 4$ $\text{nsp}_2(G_3) = 5$



G_4 : example for case 4 G_5 : example for case 5
 $\text{nsp}_2(G_4) = 10$ $\text{nsp}_2(G_5) = 6$

Figure 4: Five examples for Theorem 14.

References

- [1] G. J. Chang, S. Juan and D. Liu, *No-hole 2-distant colorings for unit interval graphs* *Ars Combinatoria*, to appear.
- [2] G. J. Chang, D. D.-F. Liu and X. Zhu, *Distance graphs and T-coloring*, *J. Comb. Theory, Series B* 75 (1999) 259-269.
- [3] M. B. Cozzens and F. S. Roberts, *T-colorings of graphs and the channel assignment problem*, *Congr. Numer.* 35 (1982) 191-208.
- [4] M. B. Cozzens and F. S. Roberts, *Greedy algorithms for T-colorings of complete graphs and the meaningfulness of conclusions about them*, *J. Comb. Inform. Syst. Sci.* 16 (1991) 286-299.

- [5] J. R. Griggs and D. D.-F. Liu, *The channel assignment problem for mutually adjacent sites*, J. Comb. Theory, Series A 68 (1994) 169-183.
- [6] W. K. Hale, *Frequency assignment: theory and applications*, Proc. IEEE 68 (1980) 1497-1514.
- [7] S. J. Hu, S. T. Juan and G. J. Chang, *T-Colorings and T-edge spans of graphs*, Graphs and Combin. 15 (1999) 295-301.
- [8] S. T. Juan, *The No-hole T-Coloring Problem*, Master Thesis, Dept. Applied Math., National Chiao Tung Univ., June 1996.
- [9] T. A. Lanfear, *Radio frequency assignment and graph coloring*, Presented at the Third Advanced Research Institute in Discrete Applied Mathematics, RUTCOR, Rutgers University, New Brunswick, NJ (May 1988).
- [10] D. D.-F. Liu, *T-colorings of graphs*, Disc. Math. 101 (1992) 203-212.
- [11] D. D.-F. Liu, *On a conjecture of T-colorings*, Congr. Numer. 103 (1994) 27-31.
- [12] D. D.-F. Liu, *T-graphs and the channel assignment problem*, Disc. Math. 161 (1996) 197-205.
- [13] D. D.-F. Liu and R. Yeh *Graph homomorphism and no-hole T-coloring*, Congressus Numerantium 138 (1999) 39-48.
- [14] J. H. Rabinowitz and V. K. Proulx, *An asymptotic approach to the channel assignment problem*, SIAM J. Alg. Disc. Math. 6 (1985) 507-518.
- [15] A. Raychaudhuri, *Further results on T-coloring and frequency assignment problem*, SIAM J. Disc. Math. 7 (1994) 605-613.
- [16] F. S. Roberts, *No-hole 2-distant colorings*, Math. Comp. Modeling 17 (1993) 139-144.
- [17] D. Sakai and C. Wang, *No-hole (r + 1)-distant colorings*, Disc. Math. 119 (1993) 175-189.
- [18] B. A. Tesman, *T-Colorings, List T-Colorings and Set T-Colorings of Graphs*, Ph.D. Dissertation, Dept. Math., Rutgers University (1989).